EXPLANATION OF PLATE II, WITH DESCRIPTION OF NEW VARIETY OF LIMENITIS URSULA.

BY H. H. NEWCOMB, BOSTON, MASS.

Fig. 1, Thecla critola, Hewitson, & Patagonia Mts., Arizona. (E. J. Oslar.)

Fig. 2, Lycaena pseudargiolus var. arizonensis, Edw., hermaphrodite; left side \circlearrowleft , right side \circlearrowleft . Huachuca Mts., Arizona. (E. J. Oslar.)

Fig. 3, Papilio polyxenes, Fabr. ♂, aberrant; lacking the inner row of yellow spots across the primaries. Albany, N. Y. (J. H. Cook.)

Fig. 4, Melitaca phaeton, Dru., ♂, aberrant; having but one row of white spots across the wings and with the marginal lunules absent. Milton, Mass. (II. II. Newcomb.)

Fig. 5, M. phaeton var. superba, Strecker, &, (under side). Milton, Mass. (W. D. Denton.) I have another from the same locality on which the white is more diffused but less distinct.

Fig. 6, Colias philodice Godt. ♀ aberrant; the primaries being melanie. Sharon, Mass. (A. C. Sampson.)

Fig. 7, Limenitis ursula var. alhofasciata, Newcomb, 3. (Previously figured, uncolored, Psyche, Vol. XI, Pl. III.) Sharon, Mass. (A. C. Sampson.) See description on next page.

Fig. 8, same as above, 2. Bedford, Mass. (L. W. Swett.)

Fig. 9, Chrysophauus hypophlaus Bdv. ♂ aberrant; left side normal, right side var. fasciata Streck. Sharon, Mass. (A. C. Sampson.)

Fig. 10, C. hypophlaeas var. fasciata $\,^{\circ}\,$, albinie. Princeton, Mass. (L. W. Swett.)

Fig. 11, C. hypophlaeas var. fasciata, A. Wellesley, Mass. (II. H. Newcomb.)

Fig. 12, Phyciodes tharos, var. packardii Saund, ♂. Sharon, Mass. (A. C. Sampson.)

Fig. 13, P. tharos, Dru. ♀ albinic. I believe this example of albinism to be unique in this species. Dorchester, Mass. (II. II. Newcomb.)

Fig. 14, Limenitis sp. ♂. This butterfly which has been previously figured in Psyche (Vol. XI, Pl. 1, fig. 4) but uncolored, may be a cross between archippus Cram, and ursula Fabr, as suggested by Field (Psyche, Vol. XI, p. 1). There are certainly some grounds for this assumption: for instance the marginal row of white dots on the upper surface is distinctly traceable to archippus, while on the under side